


Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1




3662
3663
3664
3665
3666
3667
3668
3669
3670
3671
3672
3673
3674
3675
3676
3677
3678
3679
3680
3681

T. Kumar Srinivasan. Heterologous expression of Arabidopsis NPR1 (AtNPR1) enhances oxidative stress tolerance in transgenic tobacco plants. <i>Biotechnol Lett.</i> 2009. 31:1343-51
A. Botti Balestrazzi. Expression of the PsMTA1 gene in white poplar engineered with the MAT system is associated with heavy metal tolerance and protection against 8-hydroxy-2'-deoxyguanosine mediated-DNA damage. <i>Plant Cell Rep.</i> 2009. 28:1179-92
X. Tan Wan. Increased tolerance to oxidative stress in transgenic tobacco expressing a wheat oxalate oxidase gene via induction of antioxidant enzymes is mediated by H ₂ O ₂ . <i>Physiol Plant.</i> 2009. 136:30-44
M. Miller Tunc-Ozdemir. Thiamin confers enhanced tolerance to oxidative stress in Arabidopsis. <i>Plant Physiol.</i> 2009. 151:421-32
K. Ogawa Ishikawa. Modulation of the poly(ADP-ribosyl)ation reaction via the Arabidopsis ADP-ribose/NADH pyrophosphohydrolase, AtNUDX7, is involved in the response to oxidative stress. <i>Plant Physiol.</i> 2009. 151:741-54
G. Havaux Levesque-Tremblay. The chloroplastic lipocalin AtCHL prevents lipid peroxidation and protects Arabidopsis against oxidative stress. <i>Plant J.</i> 2009. 60:691-702
L. M. Abbasi Voll. Are there Specific In Vivo Roles for alpha- and gamma-Tocopherol in Plants?. <i>Plant Signal Behav.</i> 2007. 2:486-8
A. Bonadei Balestrazzi. Leaf-associated bacteria from transgenic white poplar producing resveratrol-like compounds: isolation, molecular characterization, and evaluation of oxidative stress tolerance. <i>Can J Microbiol.</i> 2009. 55:829-40
S. G. Kim Kim. Overexpression of rice isoflavone reductase-like gene (OsIRL) confers tolerance to reactive oxygen species. <i>Physiol Plant.</i> 2010. 138:1-9
S. Misumi Hirooka. Expression of the Cyanidioschyzon merolae stromal ascorbate peroxidase in Arabidopsis thaliana enhances thermotolerance. <i>Plant Cell Rep.</i> 2009. 28:1881-93
D. Casagrande Tarantino. Knocking out of the mitochondrial AtFer4 ferritin does not alter response of Arabidopsis plants to abiotic stresses. <i>J Plant Physiol.</i> 2010. 167:453-60
R. Kim Ahmad. Simultaneous expression of choline oxidase, superoxide dismutase and ascorbate peroxidase in potato plant chloroplasts provides synergistically enhanced protection against various abiotic stresses. <i>Physiol Plant.</i> 2010. 138:520-33
F. Xu Liu. Gene expression profiles deciphering rice phenotypic variation between Nipponbare (Japonica) and 93-11 (Indica) during oxidative stress. <i>PLoS One.</i> 2010. 5:e8632
G. Gerhardt Sanchez. Salicylic acid is involved in the Nb-mediated defense responses to Potato virus X in Solanum tuberosum. <i>Mol Plant Microbe Interact.</i> 2010. 23:394-405
K. M. Holmstrom Chen. Small chloroplast-targeted DnaJ proteins are involved in optimization of photosynthetic reactions in Arabidopsis thaliana. <i>BMC Plant Biol.</i> 2010. 10:43
F. Wu Li. Overexpression of chloroplastic monodehydroascorbate reductase enhanced tolerance to temperature and methyl viologen-mediated oxidative stresses. <i>Physiol Plant.</i> 2010. 139:421-34
Z. Xiao Wang. Increased vitamin C content accompanied by an enhanced recycling pathway confers oxidative stress tolerance in Arabidopsis. <i>J Integr Plant Biol.</i> 2010. 52:400-9
X. M. Lin Xu. The Arabidopsis DJ-1a protein confers stress protection through cytosolic SOD activation. <i>J Cell Sci.</i> 2010. 123:1644-51
J. Lee Zhu. A cellulose synthase-like protein is required for osmotic stress tolerance in Arabidopsis. <i>Plant J.</i> 2010. 63:128-40
M. E. Yoshida Haque. ROS resistance in Pisum sativum cv. Alaska: the involvement of nucleoside diphosphate kinase in oxidative stress responses via the regulation of antioxidants. <i>Planta.</i> 2010. 232:367-82

In Arabidopsis, NPR1 (non-expressor of pathogenesis related genes 1, AtNPR1) functions downstream of salicylic acid (SA)
Marker-free transgenic white poplar (<i>Populus alba</i> L., cv 'Villafranca') plants, expressing the PsMT (A1) gene from <i>Pisum</i>
Hydrogen peroxide (H ₂ O ₂) plays a key role in the regulation of plant responses to various environmental stresses and
Thiamin and thiamin pyrophosphate (TPP) are well known for their important roles in human nutrition and enzyme catal
Here, we assessed modulation of the poly(ADP-ribosyl)ation (PAR) reaction by an Arabidopsis (<i>Arabidopsis thaliana</i>) ADP
Lipocalins are small ligand-binding proteins with a simple tertiary structure that gives them the ability to bind small, gene
Tocopherols belong to the Vitamin E family of amphiphilic antioxidants, together with the subfamily of tocotrienols. They
The aim of this study was the isolation and characterization of the culturable bacteria inhabiting the leaves of transgenic
Isoflavone reductase is an enzyme involved in isoflavonoid biosynthesis in plants. However, rice isoflavone reductase-like
The ability of the primitive red alga <i>Cyanidioschyzon merolae</i> to adapt to high temperatures was utilized to produce ther
Ferritins are iron-storage proteins which, in Arabidopsis, have a clear role in protection against oxidative stress. Plant fer
Plants synthesize compatible solutes such as glycinebetaine (GB) in response to abiotic stresses. To evaluate the synergis
Rice is a very important food staple that feeds more than half the world's population. Two major Asian cultivated rice (<i>O</i>
To evaluate the role of salicylic acid (SA) in Nb-mediated hypersensitive resistance to Potato virus X (PVX) avirulent strain
BACKGROUND: DnaJ proteins participate in many metabolic pathways through dynamic interactions with various compo
A tomato (<i>Lycopersicon esculentum</i> Mill.) monodehydroascorbate reductase gene (LeMDAR) was isolated. The LeMDAR-
Vitamin C (L-ascorbic acid, AsA) has important antioxidant and metabolic functions in both plants and animals. Once used
Mutations in the DJ-1 gene (also known as PARK7) cause inherited Parkinson's disease, which is characterized by neuron
Osmotic stress imposed by soil salinity and drought stress significantly affects plant growth and development, but osmot
This study investigated the reactive oxygen species (ROS) tolerance mechanism of a paraquat-resistant <i>Pisum sativum</i> lin

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant

Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1
Level 1



3682
3683
3684
3685
3686
3687
3688
3689
3690
3691
3692
3693
3694
3695
3696
3697
3698
3699
3700
3701

M. D. Kim Kim. Enhanced tolerance to methyl viologen-induced oxidative stress and high temperature in transgenic potato plants overexpressing the CuZnSOD, APX and NDPK2 genes. <i>Physiol Plant</i> . 2010. 140:153-62
Y. H. Kim Kim. Transgenic poplar expressing Arabidopsis NDPK2 enhances growth as well as oxidative stress tolerance. <i>Plant Biotechnol J</i> . 2011. 9:334-47
N. Higuchi Yokotani. A novel chloroplast protein, CEST induces tolerance to multiple environmental stresses and reduces photooxidative damage in transgenic Arabidopsis. <i>J Exp Bot</i> . 2011. 62:557-69
L. Ren Sun. An h-type thioredoxin functions in tobacco defense responses to two species of viruses and an abiotic oxidative stress. <i>Mol Plant Microbe Interact</i> . 2010. 23:1470-85
C. Liu Zhang. Overexpression of SIGMEs leads to ascorbate accumulation with enhanced oxidative stress, cold, and salt tolerance in tomato. <i>Plant Cell Rep</i> . 2011. 30:389-98
M. Boycheva Raschke. Enhanced levels of vitamin B(6) increase aerial organ size and positively affect stress tolerance in Arabidopsis. <i>Plant J</i> . 2011. 66:414-32
L. Xi Zhang. Cotton GhMPK2 is involved in multiple signaling pathways and mediates defense responses to pathogen infection and oxidative stress. <i>Febs j</i> . 2011. 278:1367-78
G. Nasar Li. Arabidopsis poly(ADP-ribose) glycohydrolase 1 is required for drought, osmotic and oxidative stress responses. <i>Plant Sci</i> . 2011. 180:283-91
B. Poage Le Martret. Tobacco chloroplast transformants expressing genes encoding dehydroascorbate reductase, glutathione reductase, and glutathione-S-transferase, exhibit altered anti-oxidant metabolism and improved abiotic stress tolerance. <i>Plant Biotechnol J</i> . 2011. 9:661-73
M. Le Martret Poage. Modification of reactive oxygen species scavenging capacity of chloroplasts through plastid transformation. <i>Plant Mol Biol</i> . 2011. 76:371-84
M. D. Kim Kim. Overexpression of 2-cysteine peroxiredoxin enhances tolerance to methyl viologen-mediated oxidative stress and high temperature in potato plants. <i>Plant Physiol Biochem</i> . 2011. 49:891-7
A. Martins Bonifacio. Role of peroxidases in the compensation of cytosolic ascorbate peroxidase knockdown in rice plants under abiotic stress. <i>Plant Cell Environ</i> . 2011. 34:1705-22
S. C. Hwang Lee. Overexpression of the pepper antimicrobial protein CaAMP1 gene regulates the oxidative stress- and disease-related proteome in Arabidopsis. <i>Planta</i> . 2011. 234:1111-25
J. H. Moon Kim. Photosynthetic capacity of Arabidopsis plants at the reproductive stage tolerates gamma irradiation. <i>J Radiat Res</i> . 2011. 52:441-9
D. Olate Laporte. Glutaredoxin GRXS13 plays a key role in protection against photooxidative stress in Arabidopsis. <i>J Exp Bot</i> . 2012. 63:503-15
Y. Chu Zhou. Overexpression of Nelumbo nucifera metallothioneins 2a and 3 enhances seed germination vigor in Arabidopsis. <i>Planta</i> . 2012. 235:523-37
K. H. Alam Kim. Overexpression of a chloroplast-localized small heat shock protein OsHSP26 confers enhanced tolerance against oxidative and heat stresses in tall fescue. <i>Biotechnol Lett</i> . 2012. 34:371-7
J. Xu Xi. Loss of AtPDR11, a plasma membrane-localized ABC transporter, confers paraquat tolerance in Arabidopsis thaliana. <i>Plant J</i> . 2012. 69:782-91
J. E. Lim Hwang. Overexpression of Arabidopsis dehydration- responsive element-binding protein 2C confers tolerance to oxidative stress. <i>Mol Cells</i> . 2012. 33:135-40
A. Yamauchi Nakai. Role of acylamino acid-releasing enzyme/oxidized protein hydrolase in sustaining homeostasis of the cytoplasmic antioxidative system. <i>Planta</i> . 2012. 236:427-36

Oxidative stress is a major threat for plants exposed to various environmental stresses. Previous studies found that trans
Nucleoside diphosphate kinase 2 (NDPK2) is known to regulate the expression of antioxidant genes in plants. Previously,
Environmental stresses are major factors in limiting plant growth and crop production. To find genes improving salt toler
Various thioredoxin (Trx) proteins have been identified in plants. However, many of the physiological roles played by the
GDP-Mannose 3',5'-epimerase (GME; EC 5.1.3.18) catalyses the conversion of GDP-D-mannose to GDP-L-galactose, an im
Vitamin B(6) is an essential nutrient in the human diet derived primarily from plant sources. While it is well established a
Mitogen-activated protein kinase (MAPK) cascades play important roles in mediating pathogen responses and reactive o
Poly(ADP-ribosyl)ation is a post-translational protein modification that plays important roles in many cellular processes i
One approach to understanding the Reactive Oxygen Species (ROS)-scavenging systems in plant stress tolerance is to ma
Reactive oxygen species (ROS), including superoxide anions, hydrogen peroxide and hydroxyl radicals are generated thro
Oxidative stress is one of the major causative factors for injury to plants exposed to environmental stresses. Plants have
Current studies, particularly in Arabidopsis, have demonstrated that mutants deficient in cytosolic ascorbate peroxidases
Proteomics facilitates our understanding of cellular processes and network functions in the plant defense response durin
The developmental stage has an influence on the overall responses of plants under biotic or abiotic stress conditions. Ho
Glutaredoxins (GRXs) belong to the antioxidant and signalling network involved in the cellular response to oxidative stres
Metallothioneins (MTs) are small, cysteine-rich and metal-binding proteins which are involved in metal homeostasis and
Small heat shock proteins are involved in stress tolerance. We previously isolated and characterized a rice cDNA clone, O
Paraquat is one of the most widely used herbicides in the world. However, no paraquat transporter has been isolated in
Dehydration-responsive element-binding proteins (DREBs) regulate plant responses to environmental stresses. In the cur
Acylamino acid-releasing enzyme/oxidized protein hydrolase (AARE/OPH) has been biochemically demonstrated to be a

Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant
Not Relevant